
Spokane Regional Clean Air Agency Air Quality Report - March 2021

The maximum daily Air Quality Index (AQI) for the month was 61 (MODERATE air quality, 24-hour average $PM_{10} = 76 \ \mu g/m^3$), recorded at Spokane-Augusta & Fiske during a windstorm on the 28^{th} . The maximum AQI for $PM_{2.5}$ was 60 (MODERATE, 24-hour average $PM_{2.5} = 16.6 \ \mu g/m^3$) recorded at Spokane Valley-Broadway & Glenn on the 13^{th} . There were 26 AQI-GOOD air quality days and 5 AQI-MODERATE air quality days in March.

Figure 1 shows the maximum daily AQIs for PM₁₀ and PM_{2.5} and Figure 2 shows daily PM_{2.5} mass concentrations across the air monitoring network in Spokane County. Tables 1 and 2 contain the maximum AQI values for each pollutant for the month and for the year to date. Table 3 summarizes the year to date daily AQIs by category. Appendix 1 of this report provides information about federal air quality standards, Appendix 2 describes the AQI, and Appendix 3 provides daily air quality data for March for all monitoring stations in the Spokane region. Current and historical air quality data can be obtained from the Washington State Department of Ecology's air monitoring data website, https://enviwa.ecology.wa.gov/home/map.

<u>Figure 1</u>: Air Quality Index (AQI) values for March 2021. The data represent the maximum AQI values across all monitoring stations within Spokane County.






Table 1: Maximum AQI values and pollutant concentrations for this reporting period.

Pollutant	AQI		Location	Date							
O_3	Ground-level ozone is monitored in Spokane County from May through September.										
PM ₁₀	61 (conc. = $76 \mu g/m^3$)	Moderate	Spokane-Augusta & Fiske	3/28							
PM _{2.5}	$60 \text{ (conc.} = 16.6 \mu\text{g/m}^3\text{)}$	Moderate	Spokane Valley – Broadway & Glenn	3/13							

Table 2: Maximum AQI values and pollutant concentrations for this year to date.

Pollutant	AQI		Location	Date						
O_3	Ground-level ozone is monitored in Spokane County from May through September.									
PM ₁₀	61 (conc. = $76 \mu g/m^3$)	Moderate	Spokane-Augusta & Fiske	3/28						
PM _{2.5}	67 (conc. = $20.5 \mu\text{g/m}^3$)	Moderate	Moderate Spokane Valley – Broadway & Glenn							

Table 3: AQI summary as of March 31, 2021

Category	Number of days in March	Number of days this year to date
Good (0-50)	26	75
Moderate (51-100)	5	15
Unhealthy for Sensitive Groups (101-150)	0	0
Unhealthy (151-200)	0	0
Very Unhealthy (201-300)	0	0
Hazardous (>300)	0	0

Appendix 1 – National Ambient Air Quality Standards

The Clean Air Act requires EPA to set National Ambient Air Quality Standards (NAAQS) for six common air pollutants, carbon monoxide (CO), lead (Pb), nitrogen dioxide (NO₂), particulate matter (PM₁₀ and PM_{2.5}), ground-level ozone (O₃) and sulfur dioxide (SO₂; Table A-1). These are known as "criteria" pollutants because the US EPA established regulatory limits to concentrations in ambient air using human health or environmentally based criteria. Carbon monoxide, particulate matter and ozone are monitored in Spokane County by the Spokane Regional Clean Air Agency (SRCAA) and the Washington State Department of Ecology (Ecology).

Table A-1: National Ambient Air Quality Standards

Pollutar [links to historical tak reviews	oles of NAAQS	Primary/ Secondary	Averaging Time	Level	Form			
Carbon Monoxide (CO)		primary	8 hours	9 ppm	Not to be exceeded more than once			
Carbon Monovide (CO)		primary	1 hour	35 ppm	per year			
Lead (Pb)		primary and secondary	Rolling 3 month period	0.15 μg/m ³ (1)	Not to be exceeded			
Nitrogen Dioxide (NO ₂)		primary	1 hour	100 ppb	98th percentile of 1-hour daily maximum concentrations, averaged over 3 years			
		primary and secondary	1 year	53 ppb (2)	Annual Mean			
Ozone (O ₃)		primary and secondary	8 hours	0.070 ppm	Annual fourth-highest daily maximum 8-hour concentration, averaged over 3 years			
		primary	1 year	12.0 µg/m ³	annual mean, averaged over 3 years			
Particle Pollution (PM)	PM _{2.5}	secondary	1 year	15.0 μg/m ³	annual mean, averaged over 3 years			
		primary and secondary	24 hours	35 μg/m ³	98th percentile, averaged over 3 years			
	PM ₁₀	primary and secondary	24 hours	$150 \mu g/m^3$	Not to be exceeded more than once per year on average over 3 years			
Sulfur Dioxide (SO ₂)		primary	1 hour	75 ppb (4)	99th percentile of 1-hour daily maximum concentrations, averaged over 3 years			
		secondary	3 hours	0.5 ppm	Not to be exceeded more than once per year			

⁽¹⁾ In areas designated nonattainment for the Pb standards prior to the promulgation of the current (2008) standards, and for which implementation plans to attain or maintain the current (2008) standards have not been submitted and approved, the previous standards (1.5 µg/m3 as a calendar quarter average) also remain in effect.

⁽²⁾ The level of the annual NO₂ standard is 0.053 ppm. It is shown here in terms of ppb for the purposes of clearer comparison to the 1-hour standard level.

⁽³⁾ Final rule signed October 1, 2015, and effective December 28, 2015. The previous (2008) O_3 standards are not revoked and remain in effect for designated areas. Additionally, some areas may have certain continuing implementation obligations under the prior revoked 1-hour (1979) and 8-hour (1997) O_3 standards.

⁽⁴⁾ The previous SO_2 standards (0.14 ppm 24-hour and 0.03 ppm annual) will additionally remain in effect in certain areas: (1) any area for which it is not yet 1 year since the effective date of designation under the current (2010) standards, and (2) any area for which an implementation plan providing for attainment of the current (2010) standard has not been submitted and approved and which is designated nonattainment under the previous SO_2 standards or is not meeting the requirements of a SIP call under the previous SO_2 standards (40 CFR 50.4(3)). A SIP call is an EPA action requiring a state to resubmit all or part of its State Implementation Plan to demonstrate attainment of the required NAAQS.

Appendix 2 – Air Quality Index

The Air Quality Index (AQI) is EPA's color-coded tool for communicating daily air quality to the public and can be calculated for any of the criteria pollutants except lead, provided monitoring data are available. An index value above 100 indicates that the concentration of a criteria pollutant exceeded the limit established in the NAAQS. Categories of the AQI are "Good" (green, 0-50), "Moderate" (yellow, 51-100), "Unhealthy for Sensitive Groups" (USG; orange, 101-150), "Unhealthy" (red, 151-200), "Very Unhealthy" (purple, 201-300) and "Hazardous" (maroon, 301-500; Table A-2).

Table A-2: Air pollutant breakpoints for the Air Quality Index.

Air Quality Index	Color Code	Index		Break	Health Effects		
Levels of Health Concern		Numerical Value	O ₃ (ppm) 8-hour	PM _{2.5} (μg/m ³) 24-hour	PM ₁₀ (μg/m ³) 24-hour	CO (ppm) 8-hour	
Good	Green	0-50	0.000-0.054	0.0-12.0	0-54	0.0-4.4	Air quality is considered satisfactory and air pollution poses little or no risk.
Moderate	Yellow	51-100	0.055-0.070	12.1-35.4	55-154	4.5-9.4	Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a very small number of people who are unusually sensitive to air pollution.
Unhealthy for Sensitive Groups	Orange	101-150	0.071-0.085	35.5-55.4	155-254	9.5-12.4	People especially sensitive to air pollution may experience health effects. The general public is not likely to be affected. An AQI in this category or above indicates that air pollution exceeds levels acceptable under federal air quality standards.
Unhealthy	Red	151-200	0.086-0.105	55.5-150.4	255-354	12.5-15.4	Everyone may begin to experience health effects; members of sensitive groups may experience more serious health effects.
Very Unhealthy	Purple	201-300	0.106-0.200	150.5-250.4	355-424	15.5-30.4	Health alert: everyone may experience more serious health effects.
Hazardous	Maroon	>300	0.201 to the Significant Harm Level* (0.600 ppm, 2 hour average)	250.5+	425+	30.5+	Health warnings of emergency conditions. The entire population is more likely to be affected.

^{*}The significant harm level (SHL) is set at a level that represents imminent and substantial endangerment to public health.

Appendix 3

Table A-3: Summary air quality data for March for air monitoring stations in Spokane County – pollutant concentration and Air Quality Index. Particulate matter mass concentration is reported as 24-hour averages in micrograms per cubic meter of air (μ g/m³). Loss of power (tripped GFCI) resulted in loss of PM_{2.5} data at Airway Heights March 6th through 8th. PM₁₀ monitoring at Spokane Valley-Broadway & Glenn was established on the 24th. BAM = Beta Attenuation Monitor, TEOM = Tapered Element Oscillating Microbalance. See Appendix 2 for information about the Air Quality Index.

Pollutant Concentration Pollutant Concentration								Air Quality Index (AQI)												
Date	PMz.5 - Airway Heights (24 hour avg, μ g/m)	PMz.5 - Colbert (24 hour avg, μ g/m)	PM2.5 - Spokane, Augusta & Fiske (24 hour avg, μg/n)	РМ:5- Spokane Valley, Broadway & Glenn (24 hour avg. µg/ $\mathring{\mathbf{m}}$	PMz.5 - Spokane, Monroe & Wellesley (24 hour avg, µg/n)	PM _{2.5} - MAXIMUM	PM10 - Turnbull NWR BAM (24 hour avg, µg/n)	PM10 - Spokane, Augusta & Fiske (24 hour avg, $\mu g/\mathring{n}$	PM10 - Spokane Valley, Broadway & Glenn (24 hour avg, μg/n)	PM10- MAXIMUM	Date	PMs.s - Airway Heights	PM2.5 - Colbert	PM2.5 - Spokane - Augusta & Fiske	PMz.s - Spokane Valley, Broadway & Glenn	PMz.s - Spokane, Monroe & Wellesley	PM10 - Turnbull NWR	PM10 - Spokane, Augusta & Fiske	PM2.5 - Spokane Valley, Broadway & Glenn	MOMIXEM 23
3/1 3/2 3/3 3/4 3/5	2.5 4.8 6.0 8.2 5.4	5.5 6.8 9.0 9.1 11.2	5.5 4.7 10.7 9.4 9.0	4.0 5.6 12.2 9.8 8.5	4.0 4.9 7.9 7.7 6.9	5.5 6.8 12.2 9.8 11.2	2.5 5.5 6.9 8 7.7	21 26 40 45 37		21 26 40 45 37	3/1 3/2 3/3 3/4 3/5	10 20 25 34 23	23 28 38 38 47	23 20 45 39 38	17 23 51 41 35	17 20 33 32 29	2 5 6 7 7	19 24 37 42 34		28 51 42 47
3/6 3/7 3/8 3/9 3/10	5.6 6.6	4.6 5.0 5.3 7.3 8.3	2.5 2.9 6.5 8.7 10.6	4.5 4.6 9.0 9.6 10.8	2.9 3.6 6.5 6.3 7.9	9.0 9.6 10.8	4.7 6.7 9.5 8.3 4.1	12 15 31 35 35		12 15 31 35 35	3/6 3/7 3/8 3/9 3/10	23 28	19 21 22 30 35	10 12 27 36 44	19 19 38 40 45	12 15 27 26 33	4 6 9 8 4	11 14 29 32 32		19 21 38 40 45
3/11 3/12 3/13 3/14 3/15	5.7 5.8 9.9 11.6 3.3	6.7 9.5 12.1 13.6 6.0	9.8 12.0 14.8 13.5 4.2	12.5 11.2 16.6 13.3 6.0	6.5 7.2 11.7 11.9 3.7	12.0 16.6 13.6 6.0	8.1	32 42 37 28 22		32 42 37 28 22	3/11 3/12 3/13 3/14 3/15	24 24 41 48 14	28 40 51 54 25	41 50 57 54 18	52 47 60 54 25	27 30 49 50 15	8 10 14 11 7	30 39 34 26 20		52 50 60 54 25
3/16 3/17 3/18 3/19 3/20	4.6 9.3 5.5 2.2 1.9	5.9 8.6 7.5 3.8 4.8	5.8 11.7 7.6 3.2 3.0	7.3 11.7 7.4 4.5 4.2	5.1 9.9 5.2 3.1 2.5		6.5 4.9	28 42 43 15 10		28 42 43 15 10	3/16 3/17 3/18 3/19 3/20		25 36 31 16 20	24 49 32 13	30 49 31 19 18	21 41 22 13 10	11 19 13 6 5	26 39 40 14 9		30 49 40 19 20
3/21 3/22 3/23 3/24 3/25	1.8 3.2 2.4 2.1 2.7	4.1 4.7 5.1 4.0 4.9	1.6 3.9 4.5 4.1 4.8	3.7 5.4 5.0 5.1 5.8	2.5 3.7 5.1 3.5 3.7	5.4 5.1 5.1 5.8	5.2	10 9 12 11 14	5.6 8.1	10 9 12 11 14	3/21 3/22 3/23 3/24 3/25	8 13 10 9 11	17 20 21 17 20	7 16 19 17 20	15 23 21 21 24	10 15 21 15 15	6 3 4 9 5	9 8 11 10 13	5 7	17 23 21 21 24 24
3/26 3/27 3/28 3/29 3/30	5.1 5.7 0.5 3.0	5.8 7.0 6.8 4.6 3.2	3.2 4.9 7.1 1.4 4.0	5.4 6.2 8.0 3.9 6.0	3.2 4.7 4.3 2.0 3.3	7.0 8.0 4.6 6.0	8.2 69.5 7.1 10.7	16 16 76 11 24	10 11.5 62.1 9.8 14.7	16 16 76 11 24	3/26 3/27 3/28 3/29 3/30	21 24 2 13	24 29 28 19 13	13 20 30 6 17	23 26 33 16 25	13 20 18 8 14	6 8 58 7 10	15 15 61 10 22	9 11 54 9 14	24 29 61 19 25
3/31 AVG MAX	4.9 4.8 11.6	3.9 6.6 13.6	7.8 6.6 14.8	8.6 7.6 16.6	5.9 6.6 13.6	7.6	10	27 27 76	18.1 17 62	27 27 76	3/31 AVG MAX	20 20 48	16 27 54	33 27 57	36 31 60	25 22 50	12 9 58	25 24 61	17 16 54	36 31 61